

Measurements of the observed cross sections for $e^+e^- \rightarrow \text{exclusive light hadrons}$ containing K_S^0 meson at $\sqrt{s} = 3.773$ and 3.650 GeV

M. Ablikim¹, J. Z. Bai¹, Y. Bai¹, Y. Ban¹¹, X. Cai¹, H. F. Chen¹⁵, H. S. Chen¹, H. X. Chen¹, J. C. Chen¹, Jin Chen¹, X. D. Chen⁵, Y. B. Chen¹, Y. P. Chu¹, Y. S. Dai¹⁷, Z. Y. Deng¹, S. X. Du¹, J. Fang¹, C. D. Fu¹⁴, C. S. Gao¹, Y. N. Gao¹⁴, S. D. Gu¹, Y. T. Gu⁴, Y. N. Guo¹, K. L. He¹, M. He¹², Y. K. Heng¹, J. Hou¹⁰, H. M. Hu¹, T. Hu¹, G. S. Huang^{1a}, X. T. Huang¹², Y. P. Huang¹, X. B. Ji¹, X. S. Jiang¹, J. B. Jiao¹², D. P. Jin¹, S. Jin¹, Y. F. Lai¹, H. B. Li¹, J. Li¹, L. Li¹, R. Y. Li¹, W. D. Li¹, W. G. Li¹, X. L. Li¹, X. N. Li¹, X. Q. Li¹⁰, Y. F. Liang¹³, H. B. Liao^{1b}, B. J. Liu¹, C. X. Liu¹, Fang Liu¹, Feng Liu⁶, H. H. Liu^{1c}, H. M. Liu¹, J. B. Liu^{1d}, J. P. Liu¹⁶, H. B. Liu⁴, J. Liu¹, R. G. Liu¹, S. Liu⁸, Z. A. Liu¹, F. Lu¹, G. R. Lu⁵, J. G. Lu¹, C. L. Luo⁹, F. C. Ma⁸, H. L. Ma¹, L. L. Ma^{1e}, Q. M. Ma¹, M. Q. A. Malik¹, Z. P. Mao¹, X. H. Mo¹, J. Nie¹, R. G. Ping¹, N. D. Qi¹, H. Qin¹, J. F. Qiu¹, G. Rong¹, X. D. Ruan⁴, L. Y. Shan¹, L. Shang¹, D. L. Shen¹, X. Y. Shen¹, H. Y. Sheng¹, H. S. Sun¹, S. S. Sun¹, Y. Z. Sun¹, Z. J. Sun¹, X. Tang¹, J. P. Tian¹⁴, G. L. Tong¹, X. Wan¹, L. Wang¹, L. L. Wang¹, L. S. Wang¹, P. Wang¹, P. L. Wang¹, W. F. Wang^{1f}, Y. F. Wang¹, Z. Wang¹, Z. Y. Wang¹, C. L. Wei¹, D. H. Wei³, Y. Weng¹, N. Wu¹, X. M. Xia¹, X. X. Xie¹, G. F. Xu¹, X. P. Xu⁶, Y. Xu¹⁰, M. L. Yan¹⁵, H. X. Yang¹, M. Yang¹, Y. X. Yang³, M. H. Ye², Y. X. Ye¹⁵, C. X. Yu¹⁰, G. W. Yu¹, C. Z. Yuan¹, Y. Yuan¹, S. L. Zang^{1g}, Y. Zeng⁷, B. X. Zhang¹, B. Y. Zhang¹, C. C. Zhang¹, D. H. Zhang¹, H. Q. Zhang¹, H. Y. Zhang¹, J. W. Zhang¹, J. Y. Zhang¹, X. Y. Zhang¹², Y. Y. Zhang¹³, Z. X. Zhang¹¹, Z. P. Zhang¹⁵, D. X. Zhao¹, J. W. Zhao¹, M. G. Zhao¹, P. P. Zhao¹, B. Zheng¹, H. Q. Zheng¹¹, J. P. Zheng¹, Z. P. Zheng¹, B. Zhong⁹, L. Zhou¹, K. J. Zhu¹, Q. M. Zhu¹, X. W. Zhu¹, Y. C. Zhu¹, Y. S. Zhu¹, Z. A. Zhu¹, Z. L. Zhu³, B. A. Zhuang¹, B. S. Zou¹

(BES Collaboration)

¹ Institute of High Energy Physics, Beijing 100049, People's Republic of China

² China Center for Advanced Science and Technology(CCAST), Beijing 100080, People's Republic of China

³ Guangxi Normal University, Guilin 541004, People's Republic of China

⁴ Guangxi University, Nanning 530004, People's Republic of China

⁵ Henan Normal University, Xinxiang 453002, People's Republic of China

⁶ Huazhong Normal University, Wuhan 430079, People's Republic of China

⁷ Hunan University, Changsha 410082, People's Republic of China

⁸ Liaoning University, Shenyang 110036, People's Republic of China

⁹ Nanjing Normal University, Nanjing 210097, People's Republic of China

¹⁰ Nankai University, Tianjin 300071, People's Republic of China

¹¹ Peking University, Beijing 100871, People's Republic of China

¹² Shandong University, Jinan 250100, People's Republic of China

¹³ Sichuan University, Chengdu 610064, People's Republic of China

¹⁴ Tsinghua University, Beijing 100084, People's Republic of China

¹⁵ University of Science and Technology of China, Hefei 230026, People's Republic of China

¹⁶ Wuhan University, Wuhan 430072, People's Republic of China

¹⁷ Zhejiang University, Hangzhou 310028, People's Republic of China

^a Current address: University of Oklahoma, Norman, Oklahoma 73019, USA

^b Current address: DAPNIA/SPP Batiment 141, CEA Saclay, 91191, Gif sur Yvette Cedex, France

^c Current address: Henan University of Science and Technology, Luoyang 471003, People's Republic of China

^d Current address: CERN, CH-1211 Geneva 23, Switzerland

^e Current address: University of Toronto, Toronto M5S 1A7, Canada

^f Current address: Laboratoire de l'Accélérateur Linéaire, Orsay, F-91898, France

^g Current address: University of Colorado, Boulder, CO 80309, USA

By analyzing the data sets of 17.3 pb^{-1} taken at $\sqrt{s} = 3.773 \text{ GeV}$ and of 6.5 pb^{-1} taken at $\sqrt{s} = 3.650 \text{ GeV}$ with the BES-II detector at the BEPC collider, we measure the observed cross sections for the exclusive light hadron final states of $K_S^0 K^- \pi^+$, $K_S^0 K^- \pi^+ \pi^0$, $K_S^0 K^- \pi^+ \pi^+ \pi^-$, $K_S^0 K^- \pi^+ \pi^+ \pi^- \pi^0$, $K_S^0 K^- \pi^+ \pi^+ \pi^+ \pi^- \pi^-$ and $K_S^0 K^- \pi^+ \pi^0 \pi^0$ produced in e^+e^- annihilation at the two energy points. We set the upper limits on the observed cross sections and the branching fractions for $\psi(3770)$ decay to these final states at 90% C.L..

I. INTRODUCTION

The $\psi(3770)$ resonance is expected to decay almost entirely into $D\bar{D}$ meson pairs since its width is almost two orders of magnitude larger than that of $\psi(3686)$ [1]. In recent years, the study of the $\psi(3770)$ non- $D\bar{D}$ decays becomes an attractive study field in the charmonium energy region due to the existing puzzle that about 38% of $\psi(3770)$ does not decay into $D\bar{D}$ meson pairs [2]. To understand the possible excess of the $\psi(3770)$ cross section relative to the $D\bar{D}$ cross section, BES and CLEO Collaborations made many efforts to study the $\psi(3770)$ non- $D\bar{D}$ decays. The CLEO Collaboration measured the $e^+e^- \rightarrow \psi(3770) \rightarrow \text{non-}D\bar{D}$ cross section to be $(-0.01 \pm 0.08^{+0.41}_{-0.30}) \text{ nb}$ [3]. While the BES Collaboration

measured the $e^+e^- \rightarrow \psi(3770) \rightarrow \text{non-}D\bar{D}$ cross section to be $(-0.01 \pm 0.08^{+0.41}_{-0.30}) \text{ nb}$ [3]. While the BES Collaboration

measured the branching fraction for $\psi(3770) \rightarrow \text{non-}D\bar{D}$ decay to be $(15 \pm 5)\%$ [4, 5, 6, 7, 8], which indicates that, contrary to what is generally expected, the $\psi(3770)$ might substantially decay into non- $D\bar{D}$ final states or there are some new structure or physics effects which may partially be responsible for the largely measured non- $D\bar{D}$ branching fraction of the $\psi(3770)$ decays [9, 10]. BES Collaboration observed the first non- $D\bar{D}$ decay mode for $\psi(3770) \rightarrow J/\psi\pi^+\pi^-$, and measured its decay branching fraction to be $\mathcal{B}[\psi(3770) \rightarrow J/\psi\pi^+\pi^-] = (0.34 \pm 0.14 \pm 0.09)\%$ [11, 12]. This was confirmed by CLEO Collaboration [13]. Latter, CLEO Collaboration observed more $\psi(3770)$ exclusive non- $D\bar{D}$ decays, $\psi(3770) \rightarrow J/\psi\pi^0\pi^0$, $J/\psi\pi^0$, $J/\psi\eta$ [13], $\gamma\chi_{cJ}(J = 0, 1, 2)$ [14, 15] and $\phi\eta$ [16], etc. Summing over these measured branching fractions yields the sum of the branching fractions for the $\psi(3770)$ exclusive non- $D\bar{D}$ decays not more than 2%. In addition, BES and CLEO Collaborations also attempted to search for other $\psi(3770)$ exclusive charmless decays [17, 18, 19, 20, 21] [16, 22, 23]. However, the existing results can not clarify the possible excess. For better understanding the origin of the possible excess, search for more $\psi(3770)$ exclusive charmless decays will be helpful.

In this Letter, we report measurements of the observed cross sections for the exclusive light hadron final states of $K_S^0 K^- \pi^+$ (Throughout the Letter, charge conjugation is implied), $K_S^0 K^- \pi^+ \pi^0$, $K_S^0 K^- \pi^+ \pi^+ \pi^-$, $K_S^0 K^- \pi^+ \pi^+ \pi^- \pi^0$, $K_S^0 K^- \pi^+ \pi^+ \pi^+ \pi^- \pi^-$ and $K_S^0 K^- \pi^+ \pi^0 \pi^0$ at the center-of-mass energies of 3.773 and 3.650 GeV with the same method as the one used in our previous works [19, 20, 21]. With the measured cross sections at the two energy points, we set the upper limits on the observed cross sections and the branching fractions for $\psi(3770)$ decay to these final states. The measurements are made by analyzing the data set of 17.3 pb^{-1} collected at $\sqrt{s} = 3.773 \text{ GeV}$ [called as the $\psi(3770)$ resonance data] and the data set of 6.5 pb^{-1} collected at $\sqrt{s} = 3.650 \text{ GeV}$ (called as the continuum data) with the BESII detector at the BEPC collider.

II. BESII DETECTOR

The BES-II is a conventional cylindrical magnetic detector that is described in detail in Refs. [24, 25]. A 12-layer vertex chamber (VC) surrounding the beryllium beam pipe provides input to the event trigger, as well as coordinate information. A forty-layer main drift chamber (MDC) located just outside the VC yields precise measurements of charged particle trajectories with a solid angle coverage of 85% of 4π ; it also provides ionization energy loss (dE/dx) measurements which are used for particle identification. Momentum resolution of $1.7\%\sqrt{1+p^2}$ (p in GeV/c) and dE/dx resolution of 8.5% for Bhabha scattering electrons are obtained for the data taken at $\sqrt{s} = 3.773 \text{ GeV}$. An array of 48 scintillation counters surrounding the MDC measures the time of flight (TOF) of charged particles with a resolution of about 180 ps for

electrons. Outside the TOF, a 12 radiation length, lead-gas barrel shower counter (BSC), operating in limited streamer mode, measures the energies of electrons and photons over 80% of the total solid angle with an energy resolution of $\sigma_E/E = 0.22/\sqrt{E}$ (E in GeV) and spatial resolutions of $\sigma_\phi = 7.9 \text{ mrad}$ and $\sigma_z = 2.3 \text{ cm}$ for electrons. A solenoidal magnet outside the BSC provides a 0.4 T magnetic field in the central tracking region of the detector. Three double-layer muon counters instrument the magnet flux return and serve to identify muons with momentum greater than 500 MeV/c . They cover 68% of the total solid angle.

III. EVENT SELECTION

In the reconstruction of the $K_S^0 K^- \pi^+$, $K_S^0 K^- \pi^+ \pi^0$, $K_S^0 K^- \pi^+ \pi^+ \pi^-$, $K_S^0 K^- \pi^+ \pi^+ \pi^- \pi^0$, $K_S^0 K^- \pi^+ \pi^+ \pi^+ \pi^- \pi^-$ and $K_S^0 K^- \pi^+ \pi^0 \pi^0$ final states, the K_S^0 and π^0 mesons are reconstructed through the decays of $K_S^0 \rightarrow \pi^+ \pi^-$ and $\pi^0 \rightarrow \gamma\gamma$.

For each candidate event, we require that at least four charged tracks are well reconstructed in the MDC with good helix fits, and the polar angle of each charged track satisfies $|\cos\theta| < 0.85$. The charged tracks (except for the K_S^0 meson reconstruction) are required to originate from the interaction region $V_{xy} < 2.0 \text{ cm}$ ($V_{xy} < 8.0 \text{ cm}$) and $|V_z| < 20.0 \text{ cm}$, where V_{xy} and $|V_z|$ are the closest approaches in the xy -plane and the z direction, respectively.

The charged particles are identified by using the dE/dx and TOF measurements, with which the combined confidence levels CL_π and CL_K for pion and kaon hypotheses are calculated. The pion and kaon candidates are required to satisfy $CL_\pi > 0.001$ and $CL_K > CL_\pi$, respectively. To reconstruct K_S^0 mesons, we require that the $\pi^+ \pi^-$ meson pairs must originate from a secondary vertex which is displaced from the event vertex at least by 4 mm in the xy -plane.

The photons are selected with the BSC measurements. The good photon candidates are required to satisfy the following criteria: the energy deposited in the BSC is greater than 50 MeV, the electromagnetic shower starts in the first 5 readout layers, the angle between the photon and the nearest charged track is greater than 22° [26, 27], and the opening angle between the cluster development direction and the photon emission direction is less than 37° [26, 27].

For each candidate event, there may be several different charged and/or neutral track combinations satisfying the above selection criteria for exclusive light hadron final states. Each combination is subjected to an energy-momentum conservation kinematic fit. For the processes containing π^0 meson in the final states, an additional constraint kinematic fit is imposed on $\pi^0 \rightarrow \gamma\gamma$. Candidates with a fit probability larger than 1% are accepted. If more than one combination satisfies the selection criteria in an event, only the combination with the longest decay

distance of the reconstructed K_S^0 mesons is retained.

To suppress the background from the $D\bar{D}$ decays, we use the double tag method [28] to remove the $D\bar{D}$ events. For example, for the $K_S^0 K^- \pi^+ \pi^0$ final state, we exclude the all possible events from $D\bar{D}$ decays by rejecting those in which the D and $\bar{D}$ mesons can be reconstructed in the decay modes of $D^- \rightarrow K_S^0 K^-$ and $D^+ \rightarrow \pi^+ \pi^0$, $D^- \rightarrow K^- \pi^0$ and $D^+ \rightarrow K_S^0 \pi^+$, $\bar{D}^0 \rightarrow K_S^0 \pi^0$ and $D^0 \rightarrow K^- \pi^+$ [28]. For the other final states, the events from $D\bar{D}$ decays are suppressed similarly. The remaining contaminations from $D\bar{D}$ decays due to particle misidentification or missing photon(s) are accounted by using Monte Carlo simulation, as discussed in Section V.

IV. DATA ANALYSIS

In the data analysis, these processes containing K_S^0 meson in the final state are studied by examining the invariant mass spectra of the $\pi^+ \pi^-$ combinations satisfying the above selection criteria for the K_S^0 meson reconstruction. The invariant masses of the $\pi^+ \pi^-$ combinations are calculated with the momentum vectors from the K_S^0 reconstruction. Figure 1 shows the resulting distribution of the invariant masses of the $\pi^+ \pi^-$ combinations from the selected candidates for the $K_S^0 K^- \pi^+$, $K_S^0 K^- \pi^+ \pi^0$, $K_S^0 K^- \pi^+ \pi^+ \pi^-$, $K_S^0 K^- \pi^+ \pi^+ \pi^- \pi^0$, $K_S^0 K^- \pi^+ \pi^+ \pi^+ \pi^- \pi^-$ and $K_S^0 K^- \pi^+ \pi^0 \pi^0$ final states. In each figure, the peak around the K_S^0 nominal mass indicates the production of $e^+ e^- \rightarrow$ exclusive light hadrons containing K_S^0 meson. Fitting the $\pi^+ \pi^-$ invariant mass spectra with a Gaussian function for the K_S^0 signal and a flat background yields the number of the events for each process observed from the $\psi(3770)$ resonance data and the continuum data. In the fit, the K_S^0 mass and its mass resolution are fixed at the values obtained by analyzing Monte Carlo samples.

V. BACKGROUND SUBTRACTION

Some other events may contribute to the selected candidate events for $e^+ e^- \rightarrow f$ (f represents exclusive light hadron final state). These include the events from J/ψ and $\psi(3686)$ decays due to ISR returns, the events from the other final states due to misidentifying a pion as a kaon or reverse, and the events from $D\bar{D}$ decays. The number N^b of these contaminations should be subtracted from the number N^{obs} of the candidates for $e^+ e^- \rightarrow f$. The estimation of them can be done based on Monte Carlo simulation. The details about the background subtraction have been described in Ref. [19]. For each background channel except $D\bar{D}$ decays, 50,000 or 100,000 Monte Carlo events are used in the background estimation. The Monte Carlo sample of each different background channel is from ten to several thousands times larger than the data in size.




Fig. 1: The $\pi^+ \pi^-$ invariant mass spectra of the candidates for the (a) $K_S^0 K^- \pi^+$, (b) $K_S^0 K^- \pi^+ \pi^0$, (c) $K_S^0 K^- \pi^+ \pi^+ \pi^-$, (d) $K_S^0 K^- \pi^+ \pi^+ \pi^- \pi^0$, (e) $K_S^0 K^- \pi^+ \pi^+ \pi^+ \pi^- \pi^-$ and (f) $K_S^0 K^- \pi^+ \pi^0 \pi^0$ final states selected from the $\psi(3770)$ resonance data (left) and the continuum data (right).

Monte Carlo study shows that the contaminations from $\psi(3770) \rightarrow J/\psi \pi^+ \pi^-$, $\psi(3770) \rightarrow J/\psi \pi^0 \pi^0$, $\psi(3770) \rightarrow J/\psi \pi^0$ and $\psi(3770) \rightarrow \gamma \chi_{cJ}$ ($J = 0, 1, 2$) can be neglected.

Even though we have removed the main contaminations from $D\bar{D}$ decays in the previous event selection (see section III), there are still some events from $D\bar{D}$ decays satisfying the selection criteria for the light hadron final states due to particle misidentification or missing photon(s). The number of these contaminations from $D\bar{D}$ decays are further removed by analyzing a Monte Carlo sample which is about forty times larger than the $\psi(3770)$ resonance data. The Monte Carlo events are generated as $e^+ e^- \rightarrow D\bar{D}$ at $\sqrt{s} = 3.773$ GeV, where the D and $\bar{D}$ mesons are set to decay into all possible final states with the branching fractions quoted from PDG [8].

Subtracting the number N^b of these contaminations from the number N^{obs} of the candidate events, we obtain the net number N^{net} of the signal events for each process. For the $K_S^0 K^- \pi^+$ and $K_S^0 K^- \pi^+ \pi^+ \pi^- \pi^-$ final states, for which only a few signal events are observed from the continuum data, we set the upper limits N^{up} on the number of the signal events at 90% C.L.. Here, we use the Feldman-Cousins method [29] and assume that the background is absent. The numbers of N^{obs} , N^b and N^{net} (or N^{up}) are summarized in the second, third and fourth columns of Tabs. I and II. For each process, the background events in the $\psi(3770)$ resonance data are dominant by $D\bar{D}$ decays and $\psi(3686)$ decays. While, there is no $D\bar{D}$ decay in the continuum data, and the $\psi(3686)$ production cross section at $\sqrt{s} = 3.650$ GeV is much less than that at $\sqrt{s} = 3.773$ GeV. So, the num-

ber of the background events in the continuum data can almost be negligible.

VI. RESULTS

A. Monte Carlo efficiency

To estimate the detection efficiency ϵ for $e^+e^- \rightarrow f$, we use a phase space generator including initial state radiation and vacuum polarization corrections [30] with $1/s$ energy dependence in cross section. Final state radiation [31] decreases the detection efficiency not more than 0.5%. Detailed analysis based on Monte Carlo simulation for the BES-II detector [32] gives the detection efficiencies for each process at $\sqrt{s} = 3.773$ and 3.650 GeV, which are summarized in the fifth columns of Tabs. I and II, where the detection efficiencies do not include the branching fractions for $K_S^0 \rightarrow \pi^+\pi^-$ and $\pi^0 \rightarrow \gamma\gamma$, $\mathcal{B}(K_S^0 \rightarrow \pi^+\pi^-)$ and $\mathcal{B}(\pi^0 \rightarrow \gamma\gamma)$.

B. Observed cross sections

Let $\mathcal{B}_{\pi^0} = \mathcal{B}(\pi^0 \rightarrow \gamma\gamma)$ for the modes of $K_S^0 K^- \pi^+ \pi^0$ and $K_S^0 K^- \pi^+ \pi^+ \pi^- \pi^0$, $\mathcal{B}_{\pi^0} = \mathcal{B}^2(\pi^0 \rightarrow \gamma\gamma)$ for the mode of $K_S^0 K^- \pi^+ \pi^0 \pi^0$ and $\mathcal{B}_{\pi^0} = 1$ for $K_S^0 K^- \pi^+$, $K_S^0 K^- \pi^+ \pi^+ \pi^-$ and $K_S^0 K^- \pi^+ \pi^+ \pi^+ \pi^- \pi^-$, where $\mathcal{B}(\pi^0 \rightarrow \gamma\gamma)$ is the branching fraction for the decay of $\pi^0 \rightarrow \gamma\gamma$, then the observed cross section for $e^+e^- \rightarrow f$ can be determined by

$$\sigma_{e^+e^- \rightarrow f} = \frac{N^{\text{net}}}{\mathcal{L} \times \epsilon \times \mathcal{B}(K_S^0 \rightarrow \pi^+\pi^-) \times \mathcal{B}_{\pi^0}}, \quad (1)$$

where $\mathcal{L}$ is the integrated luminosity of the data set, N^{net} is the number of the signal events, ϵ is the detection efficiency and $\mathcal{B}(K_S^0 \rightarrow \pi^+\pi^-)$ is the branching fraction for the decay of $K_S^0 \rightarrow \pi^+\pi^-$. Inserting these numbers in Eq. (1), we obtain the observed cross sections for each process at $\sqrt{s} = 3.773$ and 3.650 GeV. They are summarized in Tabs. I and II, where the first error is statistical and the second systematic. In the measurements of the observed cross sections, the systematic errors arise from the uncertainties in integrated luminosity of the data set (2.1% [4, 5]), photon selection (2.0% per photon), tracking efficiency (2.0% per track), particle identification (0.5% per pion or kaon), kinematic fit (1.5%), K_S^0 reconstruction (1.1% [27]), branching fractions quoted from PDG [33] (0.03% for $\mathcal{B}(\pi^0 \rightarrow \gamma\gamma)$ and 0.07% for $\mathcal{B}(K_S^0 \rightarrow \pi^+\pi^-)$), Monte Carlo modeling (6.0% [19, 20, 21]), Monte Carlo statistics (1.4% $\sim$ 4.4%), background subtraction (0.0% $\sim$ 3.0%) and fit to mass spectrum (0.4% $\sim$ 8.5%). Adding these uncertainties in quadrature yields the total systematic error Δ_{sys} for each mode at $\sqrt{s} = 3.773$ and 3.650 GeV.

The upper limit $\sigma_{e^+e^- \rightarrow f}^{\text{up}}$ on the observed cross sections for the $K_S^0 K^- \pi^+$ and $K_S^0 K^- \pi^+ \pi^+ \pi^- \pi^-$ final

states at $\sqrt{s} = 3.650$ GeV are set with Eq. (1) by substituting N^{net} with $N^{\text{up}}/(1 - \Delta_{\text{sys}})$, where N^{up} is the upper limit on the number of the signal event, and Δ_{sys} is the systematic error in the cross section measurement. Inserting the corresponding numbers in the equation, we obtain the upper limits on the observed cross sections for $e^+e^- \rightarrow K_S^0 K^- \pi^+$ and $e^+e^- \rightarrow K_S^0 K^- \pi^+ \pi^+ \pi^- \pi^-$ at $\sqrt{s} = 3.650$ GeV, which are also listed in Tab. II.

C. Upper limits on the observed cross sections and the branching fractions for $\psi(3770) \rightarrow f$

If we ignore the possible interference effects between the continuum and resonance amplitudes, and the difference of the vacuum polarization corrections at $\sqrt{s} = 3.773$ and 3.650 GeV, we can determine the observed cross section $\sigma_{\psi(3770) \rightarrow f}$ for $\psi(3770) \rightarrow f$ at $\sqrt{s} = 3.773$ GeV by comparing the observed cross sections $\sigma_{e^+e^- \rightarrow f}^{3.773 \text{ GeV}}$ and $\sigma_{e^+e^- \rightarrow f}^{3.650 \text{ GeV}}$ for $e^+e^- \rightarrow f$ measured at $\sqrt{s} = 3.773$ and 3.650 GeV, respectively. It can be given by

$$\sigma_{\psi(3770) \rightarrow f} = \sigma_{e^+e^- \rightarrow f}^{3.773 \text{ GeV}} - f_{\text{co}} \times \sigma_{e^+e^- \rightarrow f}^{3.650 \text{ GeV}}, \quad (2)$$

where $f_{\text{co}} = 3.650^2/3.773^2$ is the normalization factor to consider the $1/s$ cross section dependence. The results are summarized in the second column of Tab. III, where the first error is the statistical, the second is the independent systematic arising from the uncertainties in the Monte Carlo statistics, in the fit to the mass spectrum and in the background subtraction, and the third is the common systematic error arising from the other uncertainties as discussed in the subsection B.

The upper limit on the observed cross section $\sigma_{\psi(3770) \rightarrow f}^{\text{up}}$ for $\psi(3770) \rightarrow f$ at $\sqrt{s} = 3.773$ GeV is set by shifting the cross section by 1.64σ , where σ is the total error of the measured cross section. The results on $\sigma_{\psi(3770) \rightarrow f}^{\text{up}}$ are summarized in the third column of Tab. III.

The upper limit on the branching fraction $\mathcal{B}_{\psi(3770) \rightarrow f}^{\text{up}}$ for $\psi(3770) \rightarrow f$ is set by dividing its upper limit on the observed cross section $\sigma_{\psi(3770) \rightarrow f}^{\text{up}}$ by the observed cross section $\sigma_{\psi(3770)}^{\text{obs}} = (7.15 \pm 0.27 \pm 0.27)$ nb [19] for the $\psi(3770)$ production at $\sqrt{s} = 3.773$ GeV and a factor $(1 - \Delta\sigma_{\psi(3770)}^{\text{obs}})$, where $\Delta\sigma_{\psi(3770)}^{\text{obs}}$ is the relative error of the $\sigma_{\psi(3770)}^{\text{obs}}$. The results on $\mathcal{B}_{\psi(3770) \rightarrow f}^{\text{up}}$ are summarized in the last column of Tab. III.

VII. SUMMARY

In this Letter, we present the measurements of the observed cross sections for $K_S^0 K^- \pi^+$, $K_S^0 K^- \pi^+ \pi^0$, $K_S^0 K^- \pi^+ \pi^+ \pi^-$, $K_S^0 K^- \pi^+ \pi^+ \pi^- \pi^0$, $K_S^0 K^- \pi^+ \pi^+ \pi^+ \pi^- \pi^-$ and $K_S^0 K^- \pi^+ \pi^0 \pi^0$ produced in e^+e^- annihilation at $\sqrt{s} = 3.773$ and 3.650 GeV. These cross sections are obtained by analyzing the data sets of

17.3 pb⁻¹ taken at $\sqrt{s} = 3.773$ GeV and of 6.5 pb⁻¹ at $\sqrt{s} = 3.650$ GeV with the BES-II detector at the BEPC collider. By comparing the observed cross sections for each process measured at $\sqrt{s} = 3.773$ and 3.650 GeV, we set the upper limits on the observed cross sections and the branching fractions for $\psi(3770)$ decay to these final states at 90% C.L.. These measurements provide helpful information to understand the mechanism of the continuum light hadron production and the discrepancy between the observed cross sections for $D\bar{D}$ and $\psi(3770)$ production.

VIII. ACKNOWLEDGMENTS

The BES collaboration thanks the staff of BEPC for their hard efforts. This work is supported in part by

the National Natural Science Foundation of China under contracts Nos. 10491300, 10225524, 10225525, 10425523, the Chinese Academy of Sciences under contract No. KJ 95T-03, the 100 Talents Program of CAS under Contract Nos. U-11, U-24, U-25, the Knowledge Innovation Project of CAS under Contract Nos. U-602, U-34 (IHEP), the National Natural Science Foundation of China under Contract No. 10225522 (Tsinghua University).

-
- [1] MARK-I Collaboration, P. A. Rapidis, *et al.*, Phys. Rev. Lett. 39 (1978) 526.
 - [2] G. Rong, D. H. Zhang, J. C. Chen, hep-ex/0506051.
 - [3] CLEO Collaboration, D. Besson, *et al.*, Phys. Rev. Lett. 96 (2006) 092002.
 - [4] BES Collaboration, M. Ablikim *et al.*, Phys. Lett. B 641 (2006) 145.
 - [5] BES Collaboration, M. Ablikim *et al.*, Phys. Rev. Lett. 97 (2006) 121801.
 - [6] BES Collaboration, M. Ablikim *et al.*, Phys. Lett. B 659 (2007) 74.
 - [7] BES Collaboration, M. Ablikim *et al.*, Phys. Rev. D 76 (2007) 122002.
 - [8] Particle Data Group, 2007 partial updata for edition 2008 (URL: <http://pdg.lbl.gov>).
 - [9] BES Collaboration, M. Ablikim *et al.*, Phys. Rev. Lett. 101 (2008) 102004, arXiv: 0807.0494/hep-ex.
 - [10] BES Collaboration, M. Ablikim *et al.*, Phys. Lett. B 668 (2008) 263.
 - [11] BES Collaboration, J. Z. Bai *et al.*, HEP & NP 28(4) (2004) 325.
 - [12] BES Collaboration, M. Ablikim *et al.*, Phys. Lett. B 605 (2005) 63.
 - [13] CLEO Collaboration, N. E. Adam *et al.*, Phys. Rev. Lett. 96 (2006) 082004.
 - [14] CLEO Collaboration, T. E. Coans *et al.*, Phys. Rev. Lett. 96 (2006) 182002.
 - [15] CLEO Collaboration, B. A. Briere *et al.*, Phys. Rev. D 74 (2006) 031106.
 - [16] CLEO Collaboration, D. Cronin-Hennessy *et al.*, Phys. Rev. D 74 (2006) 012005.
 - [17] BES Collaboration, M. Ablikim *et al.*, Phys. Rev. D 70 (2004) 077101.
 - [18] BES Collaboration, M. Ablikim *et al.*, Phys. Rev. D 72 (2005) 072007.
 - [19] BES Collaboration, M. Ablikim *et al.*, Phys. Lett. B 650 (2007) 111.
 - [20] BES Collaboration, M. Ablikim *et al.*, Phys. Lett. B 656 (2007) 30.
 - [21] BES Collaboration, M. Ablikim *et al.*, Eur. Phys. J. C 52 (2007) 805.
 - [22] CLEO Collaboration, G. S. Huang *et al.*, Phys. Rev. Lett. 96 (2006) 032003.
 - [23] CLEO Collaboration, G. S. Adams *et al.*, Phys. Rev. D 73 (2006) 012002.
 - [24] BES Collaboration, J. Z. Bai *et al.*, Nucl. Instrum. Methods A 344 (1994) 319.
 - [25] BES Collaboration, J. Z. Bai *et al.*, Nucl. Instrum. Methods A 458 (2001) 627.
 - [26] BES Collaboration, M. Ablikim *et al.*, Phys. Lett. B 597 (2004) 39.
 - [27] BES Collaboration, M. Ablikim *et al.*, Phys. Lett. B 608 (2005) 24.
 - [28] BES Collaboration, M. Ablikim *et al.*, Nucl. Phys. B 727 (2005) 395.
 - [29] G. J. Feldman and R. D. Cousins, Phys. Rev. D 57 (1998) 3873.
 - [30] E. A. Kuraev and V. S. Fadin, Yad. Fiz. 41 (1985) 377.
 - [31] E. Barberio, Z. Was, Comput. Phys. Commun. 79 (1994) 291.
 - [32] BES Collaboration, M. Ablikim *et al.*, Nucl. Instrum. Methods A 552 (2005) 344.
 - [33] Particle Data Group, W.-M. Yao, *et al.*, J. Phys. G 33 (2006) 1.

Tab. I: The observed cross sections for $e^+e^- \rightarrow$ exclusive light hadrons at $\sqrt{s} = 3.773$ GeV, where N^{obs} is the number of events observed from the $\psi(3770)$ resonance data, N^{b} is the total number of background events, N^{net} is the number of the signal events, ϵ is the detection efficiency, Δ_{sys} is the relative systematic error of the observed cross section, and σ is the observed cross section.

$e^+e^- \rightarrow$	N^{obs}	N^{b}	N^{net}	$\epsilon(\%)$	$\Delta_{\text{sys}}(\%)$	$\sigma^{\text{obs}}[\text{pb}]$
$K_S^0 K^- \pi^+$	18.4 ± 4.6	0.1 ± 0.0	18.3 ± 4.6	10.02 ± 0.14	10.7	$15.2 \pm 3.8 \pm 1.6$
$K_S^0 K^- \pi^+ \pi^0$	41.2 ± 6.6	1.1 ± 0.2	40.1 ± 6.6	3.52 ± 0.08	11.6	$96.2 \pm 15.9 \pm 11.1$
$K_S^0 K^- \pi^+ \pi^+ \pi^-$	40.0 ± 6.5	1.0 ± 0.2	38.9 ± 6.5	3.56 ± 0.06	14.2	$91.5 \pm 15.3 \pm 13.0$
$K_S^0 K^- \pi^+ \pi^+ \pi^- \pi^0$	24.5 ± 5.2	1.5 ± 0.3	23.0 ± 5.2	0.77 ± 0.03	15.2	$253.0 \pm 57.1 \pm 38.4$
$K_S^0 K^- \pi^+ \pi^+ \pi^+ \pi^- \pi^-$	4.8 ± 2.2	0.3 ± 0.1	4.5 ± 2.2	0.84 ± 0.03	18.4	$44.4 \pm 21.9 \pm 8.2$
$K_S^0 K^- \pi^+ \pi^0 \pi^0$	19.8 ± 4.9	2.8 ± 0.5	17.0 ± 4.9	0.99 ± 0.04	14.3	$147.0 \pm 42.4 \pm 21.0$

Tab. II: The observed cross sections for $e^+e^- \rightarrow$ exclusive light hadrons at $\sqrt{s} = 3.650$ GeV, where N^{obs} is the number of events observed from the continuum data, N^{up} is the upper limit on the number of the signal events, σ^{up} is the upper limit on the observed cross section set at 90% C.L., and the definitions of the other symbols are the same as those in Tab. I.

$e^+e^- \rightarrow$	N^{obs}	N^{b}	$N^{\text{net}}(\text{or } N^{\text{up}})$	$\epsilon(\%)$	$\Delta_{\text{sys}}(\%)$	$\sigma^{\text{obs}}(\sigma^{\text{up}})[\text{pb}]$
$K_S^0 K^- \pi^+$	2	0.0	< 5.91	10.55 ± 0.15	12.9	< 14.3
$K_S^0 K^- \pi^+ \pi^0$	7.7 ± 2.9	0.0	7.7 ± 2.9	3.62 ± 0.09	11.6	$47.9 \pm 18.0 \pm 5.6$
$K_S^0 K^- \pi^+ \pi^+ \pi^-$	13.4 ± 3.8	0.0	13.4 ± 3.8	3.66 ± 0.06	14.1	$81.4 \pm 23.1 \pm 11.5$
$K_S^0 K^- \pi^+ \pi^+ \pi^- \pi^0$	4.6 ± 2.5	0.0	4.6 ± 2.5	0.87 ± 0.03	17.2	$119.0 \pm 64.7 \pm 20.5$
$K_S^0 K^- \pi^+ \pi^+ \pi^+ \pi^- \pi^-$	0	0.0	< 2.44	0.95 ± 0.03	18.1	< 69.7
$K_S^0 K^- \pi^+ \pi^0 \pi^0$	3.3 ± 2.0	0.0	3.3 ± 2.0	1.12 ± 0.05	14.1	$67.1 \pm 40.7 \pm 9.5$

Tab. III: The upper limits on the observed cross section $\sigma_{\psi(3770) \rightarrow f}^{\text{up}}$ at $\sqrt{s} = 3.773$ GeV and the branching fraction $\mathcal{B}_{\psi(3770) \rightarrow f}^{\text{up}}$ for $\psi(3770) \rightarrow f$ are set at 90% C.L.. The $\sigma_{\psi(3770) \rightarrow f}$ is calculated with Eq. (2), where the first error is the statistical, the second is the independent systematic, and the third is the common systematic error. The upper * denotes that we neglect the contributions from the continuum production.

Decay Mode	$\sigma_{\psi(3770) \rightarrow f}(\text{pb})$	$\sigma_{\psi(3770) \rightarrow f}^{\text{up}}(\text{pb})$	$\mathcal{B}_{\psi(3770) \rightarrow f}^{\text{up}}$
$K_S^0 K^- \pi^+$	$15.2 \pm 3.8 \pm 0.2 \pm 1.6^*$	< 22.0	3.2×10^{-3}
$K_S^0 K^- \pi^+ \pi^0$	$51.4 \pm 23.2 \pm 2.6 \pm 5.8$	< 90.7	13.3×10^{-3}
$K_S^0 K^- \pi^+ \pi^+ \pi^-$	$15.3 \pm 26.5 \pm 2.2 \pm 2.1$	< 59.0	8.7×10^{-3}
$K_S^0 K^- \pi^+ \pi^+ \pi^- \pi^0$	$141.6 \pm 83.2 \pm 14.7 \pm 20.7$	< 284.3	41.8×10^{-3}
$K_S^0 K^- \pi^+ \pi^+ \pi^+ \pi^- \pi^-$	$44.4 \pm 21.9 \pm 2.1 \pm 7.9^*$	< 82.7	12.2×10^{-3}
$K_S^0 K^- \pi^+ \pi^0 \pi^0$	$84.2 \pm 57.2 \pm 8.3 \pm 11.2$	< 180.4	26.5×10^{-3}